ACFEL - TR- 75

CORPS OF ENGINEERS, U.S. ARMY

10 10 10

(227E)



DESCRIPTION AND CLASSIFICATION OF FROZEN SOILS

Based Upon a Joint Effort of the Division of Building Research, National Research Council of Canada, and the Arctic Construction and Frost Effects Laboratory, U. S. Army Corps of Engineers

Ġ

DRAFT



TECHNICAL REPORT NO. 75

Arctic Construction and Frost Effects Laboratory
U.S. Army Engineer Division, New England
Waltham, Massachusetts


for

Office of the Chief of Engineers
Civil Engineering Branch
Engineering Division
Military Construction



January 1961


DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

CORPS OF ENGINEERS, U.S. ARMY

DESCRIPTION AND CLASSIFICATION OF FROZEN SOILS



TECHNICAL REPORT NO. 75


Arctic Construction and Frost Effects Laboratory
U.S. Army Engineer Division, New England
Waltham, Massachusetts

for

Office of the Chief of Engineers
Civil Engineering Branch
Engineering Division
Military Construction

Details of Illustrations in this document may be belter tudied on microfiche

January 1961



PREFACE

The heart of the description and classification of frozen soils shown in columns (1) through (6) of Figure 2 of this report represents the joint efforts of representatives of the Building Research Division, National Research Council of Canada, and of the Arctic Construction and Frost Effects Laboratory, U. S. Army Engineer Division, New England. It is based on the experience of these organizations over several years with various forms of a system originally devised by the Arctic Construction and Frost Effects Laboratory in 1952.* The remainder of Figure 2, and of the report, is a contribution of the Arctic Construction and Frost Effects Laboratory.

This presentation is the product of a program of studies being conducted for the Chief of Engineers, Department of the Army, under the administrative direction of the Civil Engineering Branch, Engineering Division, Military Construction. The program is aimed at developing engineering criteria for design and construction in arctic and subarctic regions and in areas of seasonal frost.

[&]quot;Published as Appendix A of Vol. 1 of "Investigation of Description, Classification and Strength Properties of Frozen Soils," by Arctic Construction and Frost Effects Laboratory, issued as Report 8 of U. S. Army Snow, Ice and Permafrost Research Establishment, June 1952.

CONTENTS

		Page
Preface		* ii
Synopsis .	· · · · · · · · · · · · · · · · · · ·	iv
Introduction	• • • • • • • • • • • • • • • • • • • •	1
Features of t	he Frozen Soil Classification System	2
Ice or Water	Content of Frozen Saturated Soils	7
Graphical Pr	esentation of Soils Data	8
	ILLUSTRATIONS	
Figure 1 Illu	stration of Frozen Soil Terminology	
Figure 2 Des	scription and Classification of Frozen Soils	
Figure 3 Uni	ified Soil Classification	
Figure 4 Pho	otographs of Frozen Soil Types	
Figure 5 Pho	otographs of Frozen Soil Types	
	l Dry Unit Weight, Ice Volume, and Water Content lationships	
	strative Example of Frozen Soil Classification	

SYNOPSIS

The description and classification of frozen soils presented herein is an extension of the Unified Soil Classification System adopted by the U. S. Army Corps of Engineers and the U. S. Eureau of Reclamation in 1952.

Descriptions, based on physical appearance, are non-genetic and are applicable to both naturally and artificially frozen soils. Field identification data pertaining to frozen soils and those pertinent properties of frozen materials which can be measured by physical tests are indicated. Also, guides are presented for construction on soils subject to freezing and thawing. The report includes photographic illustrations of frozen soil types; a chart showing relationships between unit dry weight of soil, water content, and ice volume; and an illustrative example of graphical presentation of frozen soil data.

INTRODUCTION

1. When the Unified Soil Classification System* is extended to classification of frozen soils, special expansion of the system is required in order to meet engineering and scientific needs for adequate and concise identification of the materials. Identification of seasonally frozen soil or permafrost according to structural divisions caused by freezing and thawing such as "suprapermafrost" or "annual frost zone," illustrated in Figure 1, provides no information on those factors of appearance and physical properties which are essential guides to the nature and behavior of the materials in the frozen state and to the changes which may occur upon thawing. Also, such identification is not applicable to specimens frozen in the laboratory. Therefore, a frozen soil description and classification system, which is independent of the geologic history or mode of origin of the material, is needed. This system should also be capable of easy expansion or contraction in order to provide any desired degree of detail. The system described herein affords these characteristics.

^{*}Described in Technical Memorandum No. 3-357, U. S. Army Waterways Experiment Station, March 1953, with Appendixes A and B.

FEATURES OF THE FROZEN SOIL CLASSIFICATION SYSTEM

- 2. Parts of the System. The system for describing and classifying frozen soil is shown in Figure 2. As indicated in the first column of Figure 2, the frozen soil is identified in three steps, denoted as Parts I, II, and III. Under Part I the soil phase is identified independently of the frozen state; the Unified Soil Classification System is used, a summary of which is shown in Figure 3. Under Part II, the soil characteristics resulting from the frozen state of the material are added to the soil description. Under Part III, important ice strata found in the soil are described.
- 3. Classification of Frozen Soil Major Groups. As shown in columns (2) and (3) of Figure 2, under Part II, frozen so sare divided into two major groups: soils in which segregated ice is not visible to the unaided eye (designation N), and soils in which segregated ice is visible (designation V). Since, as will be described below, ice layers exceeding 1 inch in thickness are identified separately, the latter major grouping is applied only to soil containing ice layers 1 inch or less in thickness.
- 4. Frozen soils in the N group will commonly, on inspection by the unaided eye, reveal the presence of ice within the soil voids by crystalline reflections or by a sheen on fractured or trimmed surfaces; however, the appearance is given that the water has frozen within the original voids in the soil, without segregation. Frozen soils in the V group give the opposite impression, and segregated ice is visible not morely as pin point crystalline reflections or a diffuse sheen but as separate ice inclusions of measureable dimensions.

5. Frozen Sons in which Segregated Ice is not Visible. As shown in columns (4) and (5) of Figure 2, materials in which segregated ice is not visible to the unaided eye (designation N) are divided into two types:

Nf (ne non-resuble friable). This is poorly bonded or friable material in which segregated he is not visible to the unaided eye. This condition exists when the degree of saturation is low. This type of frozen soil is illustrated in photographs 1 and 2 of Figure 4.

No (ice ron visible; bonded). This is well bonded frozen soil in which the ice cements the material into a hard solid mass, but segregated ice is not visible to the unaided eye. Soils showing this characteristic are generally at a moderate to high degree of saturation. When at high degree of saturation, they may or may not contain substantial quantities of microscopic segregated ice. On basis of detailed examinations and tests this sub-group may be further divided into the following sub-categories:

No segregated ice is present, either visible to the unaided eye or mirroscopic. This type of frozen soil is illustrated in photographs 1 and 3, Figure 4.

Nhe (contains excess ice, microscopic). This condition may occur in very fine silty sands or coarse silts where excess ice is present but is so uniformly distributed that it is not readily apparent to the unaided eye.

Appreciable settlement may occur in such soils upon thawing. This type of frozen soil is illustrated in photograph 4, Figure 4.

segretied the isometric the unaided eye (designation V) are divided into the tollowing tour sub-groups, arranged approximately in sequence of increasing the court of its sound of its sound of the court of its sound of its soun

 ∇x . $e^{-is} > e^{-is} div.d$. $e^{-is} ysths \rightarrow r$ inclusions)

 V_{1} (i.e. v.s.b.e. ice cost. gs on particles)

Vr lice v.s.ble rand m / irregularly oriented ice formations)

Vs the visible; stratified or distinctly oriented ice formations)

The Ve type of frozen some shown in photograph 5, Figure 4; Vr types of frozen some are insustrated in photographs 6 and 7, Figure 5, and Vs types in photographs 8, 9, and 10, Figure 5.

7. Description of Substantia. Ice Strata. Referring to columns (2) and (3) of Figure 2 under Part III, substantial ice strata greater than 1 inch in thickness are designated separately as ICE. As shown in columns (4) and (5) of Figure 2, the identification may fall into either of the following two broad categories:

Ire (ire without soil inclusions)

8. Identification and Description. Field identification guidance is presented in column (6) of Figure 2. In addition to determination of major group and sub-group in accordance with columns (2) through (5) of Figure 2, additional descriptive terms and data may be used as indicated therein.

Some of the sol's found in permafrost regions may also be described in exploration logs by special terms (such as "muskeg") for additional clarification.

- 9. When more than one sub-group transcreristic is present in the same material, multiple sub-group designations may be used, as Vs, r. Photograph 2, Figure 4, shows an example of frozen soil of the latter type.
- 10. When greater detail and more specific information is desired than is obtainable from visua. Inspection, physical tests and measurements may be performed on the frozen soil as indicated in column (7) of Figure 2. A camera, a small-power hand magnifying lens, and pint, size graduated jars should be standard items of field equipment for soil and survey crews. To obtain a rough estimate of the possible presence of excess ice, a simple field test can be made by placing a lump of frozen soil in a jar, allowing it to melt and visually observing the relative volume of supernatant or free water standing above the soil after the lump has melted. By initially performing this test with specimens of known ice content, a basis for field judgement can be established. Since proportions of ice and soil may vary widely, it may sometimes be difficult to decide without such a test whether a given material falls, for example, in the category of frozen soil or of ice with soil inclusions. Material containing as much as 80 percent of ice by volume and only 20 percent soil can sometimes give the appearance of being mostly soil. When more exact evaluation of presence of excess ice is required, specimens may be thawed in the laboratory in consolidometers or rubber membranes, or material may be thawed in place in the field.

- 11. Only needed portions of the detail and descriptive material outlined in columns (4) through (7) of Figure 2 should be used. In many of the simpler engineering applications, only a few of the most important elements need be recorded. For some investigations it may be satisfactory to use the Nb designation without breakdown into Nbn or Nbe categories. In other applications it might even be sufficient to use only the N and V major group designations, to indicate whether or not segregated ice is visible. On the other hand, in many scientific studies very detailed records may be necessary.
- In a Characteristics. For engineering purposes, it is of very great importance to known whether significant settlement will take place upon thawing of the frozen soil. If the amount of ice present will produce more water upon melting than can be held in the veids of the soil, then the material is thaw-unstable to a degree that is dependent upon the amount of the excess ice and the soil density. If a , the melt water can be absorbed by the soil voids without significant settlement, then the soil can be considered thaw-stable.

 Columns (8) and (9) of Figure 2 present guides for construction on soils subject to freezing and thawing. The thaw characteristics shown in column (8) are particle and yellowing. Freezen soils designated as Nf and Nbn are usually thaw stable, that is, no detrimental settlement of structures would normally be anticipated if thawing accurred. Frozen soils in all other sub-groups are potentially thaw unstable and significant settlement of structures founded thereon may occurred.

- 13. Frozen open-work gravel is a special type of material which often proves difficult to evaluate as to its thaw-settlement potential. Although substantial amounts of pure ice are apparent in the voids of such material, sufficient point contacts between particles may exist to limit settlement on thaw to minor amounts. In critical cases, field thaw-settlement tests, using loaded plates and steam thawing, may be necessary.
- 14. Frozen bedrock does not always provide a thaw-safe foundation.

 Therefore, when bedrock is encountered in subfreezing temperatures, careful observations should be made to determine the quantity and mode of occurence of all ice formations in bedding planes, fissures, or other spaces.


ICE OR WATER CONTENT OF FROZEN SATURATED SOILS

15. In considerations involving frozen soils, the generally prevailing conditions include complete saturation of the soil phase and all of the water frozen. For these conditions, and assuming a specific gravity of the soil particles of 2.70, the relationships between the unit dry weight of soil, water content, and ice volume are shown in Figure 6. This chart may be used by designers or field engineers for rapid estimation of the relationships between these variables. Use of the chart is indicated by the following example and illustrated by lines and arrows on Figure 6. Assume a specimen of frozen silt with excess ice estimated at approximately 60 percent. Based on the appearance of the silt layers in the core, it is estimated that the normal dry unit weight of the silt is fairly high, say 95 pcf. The chart is then entered at 95 pcf on the left and a horizontal line is extended to the

intersection of the sloping 60 percent excess ice line. The total porosity, n, which in this case equals the proportion of ice volume of the total specimen, is then observed on the scale at the bottom of the plot (77 percent). The intersection of the vertical line (77 percent porosity) with the 100 percent saturation line indicates on the left-side scale the equivalent overall dry unit weight of the frozen specimen, i.e., 38 pcf. The curve in Figure 6 marked "Percent Volume of Ice vs Water Content" shows the relationship between the water content of a frozen specimen and total volume of ice or porosity, n. For a porosity of 77 percent in the above example, the water content indicated by the right-side scale would be approximately 114 percent.

GRAPHICAL PRESENTATION OF SOILS DATA

drawings as schematic representations of the borings or test pits, with the various soils encountered shown by appropriate symbols. The recommended procedure for graphical presentation of frozen soil classification consists of showing the applicable letter symbols for the soil phase in accordance with the Unified Soi! Classification System for unfrozen soils, followed by the frozen soil designation. An illustrative example of the use of the frozen soil classification system in a typical exploration log is shown in Figure 7. For the purpose of readily identifying the frozen soil zones, a wide line is drawn down the left side of the graphic log of the exploration within the range that the frozen material occurs.



Definitions* of Soil and Other Terms Relating to Frozen Ground Areas

Annual frost zone factive layer). The top layer of ground subject to annual freezing and thawing. In arctic and subarctic regions where annual freezing penetrates to the permafrost table, suprapermafrost and the annual frost zone are identical.

Excess ice. Ice in excess of the fraction which would be retained as wat r in the soil voids upon thawing.

Frost table. The surface, usually irregular, which represents the penetration, at any time in spring and summer, of thawing of the seasonal frozen ground.

Frozen zone. A range of depth within which the soil is frozen. The frozen zone may be bound both top and bottom by unfrozen soil, or at the top by the ground surface.

Ground ice. A body of more or less clear ice within frozen ground,

lee wedge. A wedge-shaped ice mass in permafrost, usually associated with fissure polygons.

 $\underline{lcing},\ A$ surface recomass formed by freezing of successive sheets of water.

Muskeg. Poorly drained organic terrain consisting of a mat of vegetation overlying peat of varying thickness, from a few inches to many feet,

Permafrost. Perenially frozen ground.

Permafrost table. The surface which represents the upper limit of

Pereletok. A frozen layer at the base of the active layer which remains unthawed for one or two summers.

Residual than zone. A layer of unfrozen ground between the permafrost and the annual trost zone. This layer does not exist where annual frost extends to permafrost.

Suprapermafrost. The entire layer of ground above the permafrost table.

^{*}For more complete list of definitions, see Hennion, F., Frost and Permafrost Definitions - High ery Research Board Bulletin 114, 1988,

DESCRIPTION AND CLASSIFICATION OF FROZEN SOILS

2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2					,	Class, by Soil Prase 29 the Unit	Unit ed Soul Class had on System		
		***	ensure single	6			Berthert Properties of Frozer Marks Als Will Marks Medals	3	عد بادر ومداد در در ود در ود در
		Des. 164*100	10 12 18.1	Des gration	-	field tdentification	to Pres Lat Tests to Supplement	\$5. 1578.09.2)	
	•	-		53		16,	1,	, o	
	9 P	-	Facely banded	ž	Ident to Sr would eight	anestad avietatio o' noite deserve and animal of deserve animal of deserve animal of the serve animal of t		3	The potential and the state of
		•	ani second second ani ani ani	2	nand regn 'ern, en' av Naturaled 'estreat's deg Loa 'estre proposition's around larger particles.	And sept to the term of the te	Dens is and so a ser's		
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1			Individual rec crystals or reclesions	\$	bor ice prase, record	prase, record the following as applicable:	Street 31 - Corporation of Corporati		A TOTAL TO THE PROPERTY OF THE
(2) (3) (4) (4) (5) (7) (7) (7) (7)	# A		ice coatings on particles	٨٠	Location Discretation Thickness Lend	Size Shape Pattern of arrangement	Elastic Properties Plastic Properties		Control of the contro
		>	.andor of .rrmq.lariy priented ic.	\$		per Part III selo-	Ice Cristal Structure [using optical instruments] a Orientation Asis b, Cristal Share c, Cristal	4, 49 - 1, 49 - 1, 49 - 1, 50 - 1, 50	Service and the service of the servi
			Stratified or Sistinctly Sirented ice formations	*	Estimate volume of vising percent of total sample	sample segregated ice present 43 sample solume.	i i		Section 1 and 1 an
.27	901 901 901 901 901 901		ice att soil	soil type		re ter from			
Seastan 100	11.00-05.5		ice mithout soil inclusions	9 21	THE CLEAK SOFT CLOUDY (of mays, not POROUS sociated CANDLES crystals) GRANLLAR STRAILIED	COLO-LESS CONTENS FEW SAAT THE SLUT SLUT SLUT SLUT SLUT SLUT SLUT SLUT	Cracing (Crescal		
ice Coatings on soil particles of crystals, which	Particles at	e discernible 0+1 mass. The	(14 - 17) Section 1985. The control of the control of the control of the control of the larger of the control	on or below lociated with ig action.		ry-bonged symmittes that the s first the fragen soil consequent Lite denotes ratherely mode bon en up.	Populariones in writes that the Soil satisfactor are wally held tontier in the soil manifest for the soil manifest for the soil consequency. That the finance of changes and consequency, the soil decided decided and soil of the soil for the soil particles. Material is easily found to	1 0: rearing.	The state whom a second control of the secon
100 7093031 12		on a process of the	east for a law of any serior of the solution of the state of a solution of the serior of a solutions.	offer ice		-Stable frozer son s do not, o	Tramistable frozer son s oo not, or traming, show loss of strength seio-rosmal, long-time trames walkes not produce betrimental settlement.	Velo- 101981.	10 - 10 - 10 - 10 - 10 - 10 - 10 - 10 -
Cloudy Ice 13 re	ansparent and elataely opas ally sound at	dur due to ent	Liber . I strangiver and unitary only a moderate number of air bubbles. Cloudy (ce is refaired) power due to entrained air bubbles or other reasons. Arich is essertally sound and nonneptivious. (6)	of air tutbl	(e) but	-Unstable tratem soils stom, or ral, long-time tramps values and the melting of the excess (ce.)	Inau Unstable frien soils stok, on than no is princant loss of strenot belowering the mornal, long-time transcialists and/or significant intlement, as a direct from of the melting of the entess its in the foot.	renath Below Lourest result	The state of the s
Preparation of the state of the	A TO THE STATE OF	ong crestal in row the freezy	brough the ontoins mimics social whalip represented individually resulting from after my usually resulting from the my of my and my selected that or other water is not a first or other forces, the	nd usually re mac of sait :		Fren rock is encountered, stan	NOTES: (a) Free rook is encountered, standard for classify after terminations should be used. (b) Free rook is encountered, standard for classify after termination increases of the	ny steuka de use	Amenis pintos presentados provinces destrucios destrucios de definitarios de destrucios de de destrucios de destrucios de destrucios de destrucios de destru
**************************************	\$0000000000000000000000000000000000000	rottes or of	offs for a second control of the second seco	וייים בטוייהו		and the voids of the material by Comed surfaces. Nowever, the Imiter occupies space in each rue of frazer soils in the figure	within the condist the "attential by crystalline reflections or by a steem on tractified or frozen strates." In the complexity of the changes eye is that now of the frozen water occupies space in energy of the original voids in the 3001. "Peophosite is true of frozen soils in the 1 aroup.	non tractured or at none of the soll. The oppos	
Sonded Coperior	composed of enticular co eralls openal		dambara (co. 1) composes st. Galse, more or cas equicimensoral, ice crisions and composed software constitutions and composed software com	mas, ice cry isertially pa	100	when visual methods way be in page of excess ice can 're wade " if to melt and observing the Sime."	(c) when visual methods may be inaccounte, a simple field test to tid evaluation of volume of ercess ice can in made in placing some frozen soil in a position, allow- nion at to melt and observing the quantity of supermatant mater as a percent of total volume.	rali ar, allom recent of tota	
identify, comp	a the groat	and see as de	ispers, ice acception in the growth of ice as distinct lenses, layers, weins, you wasses the world, commons, with or always privated normal to direction of heal loss.	ers, veras, b		Where special forms of ice, s	(a) where special forms of ice, such as nearfrost, can be distinguished. Fore everyleist description should be given.	137ec. 30re es-	
Section Section 1998		5 201 2 2 2 2 1 1 2 2 2 2 1 1 1 1 2 2 2 2	recits once with the that is a suit particles are strongly held together by the ice and that the requirement of the second that he region soil possesses relatively high resistance to chipping of breaking.	eld together tochippings		Observer should be careful to	(e) Observer should be careful to aword being migled by surface scratting or frost colling on the ice.	ratifies or frost	***************************************

	Laboratory Classification	-	part.		58, 58, 56, 56, 56, 56, 56, 56, 56, 56, 56, 56	hand how how the state of the s	forms is a section of the section of	to segant of segants of section of section	tree under the percent and the	Attended late above "A" line with FI greater than 7	201) sc	dž patv		S S	CHOICH LL	E AVALUE	30 MINISTER NO. 1	% or	PLANTICET CAANT POT LABORATORY CLASSIFICATION OF FINANCE AND LABORATORY CLASSIFICATION OF FINANCE AND LABORATORY CHARGES.	th tlay binder. (2) All sieve sieve on this chart are U. S. standard.	iffcation purposes,	Toughares (consistency near plantic limit)	After particles larger than the Bo. We item these remaining of parts. If to the water mark he added not it will be added to the consistency of parts. If to this large and allowed precises beautifully provided out in a trailed out by larger and allowed to lose town soliture by exponention. Then the operiment is relied out by hand on a smooth early one between the pains into a thread about one rightly inch is disserter. The thread is then folded and swolled repeatedly. During the samplakeing the soliture content is probable, when the place it is probable. After the thread cruebles, the pieces should be larged tapether and a slight mostly action continued until the laws cruebles. The thread soliture is the trailed to the place in the brightly cruebles, the wars potential and quick layer of earlier language to the brightly cruebles, the ways performed in the solitude carbon of the brightly cruebles, the ways performed in the solitudes either the compare of the brightly cruebles, the ways peat and organic fairty with occur below the Alian materials are and as a bright larger to the place of coherence of the larger alian and organic fairty with occur below the Alian Materials have a way west and appair feel at the place in late.
UTICATION and Description)	Information Department for Describing Bolia	9	For undigracted soils add information on structforthing, degree of convect- ment. constitution, maintent condition	characteri	Cive typical name; indicate approximate.	percentages of send and gravel, maxi- sam sits, angularity, surface condi- tion, and hardness of the coarse grains; local or gradings uses and	other pertinent oscinitive through	i e	Billy said, gravelly; about, 305 bard, supplier gravel particles 1/2-14. maximum sites; rounded and submittains send graves, course to fine; about 1/9 send graves.	well compacted and moist in place; al- lurial mond; (SN).			For undisturbed solls add information on structure, stratification, con-	molded states, moleture and drain-	Olve typical mase; indicate degree and	character of planticity; smouth and marines also of coarse grains; color in wer condition; odor, if any local or peologic name and other partiment	to parentheese.	Charge all, brown; alightly plastic;	numerous vertical rook boise; first and dry in place; losse; (ML).	ated by combinations of group symbols. For sample 08-00, will-graded gravel-sand mixture with flay combinations of parents and presented the presented that presented the presented that presented the presented that the presented the presented that the presented that the presented	be partorned on the base in a particles, agreemently 1/4 in. Br. field classification purposes in the other supply reserve by hand the coarse particles that inferiors with the traction.		Considerated practical Agent Laboratory. Miles the part to dry completely by considerated of party, adding where if messenty. Miles the part to dry completely by seem, was, or Liderfule, and then set it is strength by breaking and creatifully present, was, or Liderfule, and then set it is strength in a measure of the character and quantity of the reclaimed has frageria. This strength is a measure of the character and quantity of the recreasing abstracts, and the character and quantity of the recreasing abstracts. Indeed, or strength is characteristic for being of the off proper assets and the sent the measurement of the character and character and the measurement of the characteristic for being a the off the characteristic for being at the distinguished by the free has the smooth feel of filter. The measurement of the characteristic forms and fattinguished by the free has the smooth feel of filter.
GENERAL CAMEDICATION OF (Including libertifies)	Plaid Identification Procedures (Excluding particles larger than 3 in. and baing Fractions on estimated weights)		its grain since and executial of all intermediate particle since.	unge of stree with ting.	implantic flame or flame with low planticity (for identification procedure see ML below).		e compt e	mage of stress adssing.	implants fine or fine with low planticity (for identification procedure see ML below).	idestification procedures	{	(Consistency near 71.)	and R	, pedia	Biton	Silger to	THE STATE OF THE S	alignt to	ador, spongr feel strum.	gradule. For example we will	to stave also partic	rteristics)	they within the for the state of the country of the state
T.	partitionion partition larger fractions on ev	_	prais eime and il infermediate	fredominatily one sine of a runge some intermediate sines mineling.	s or flass with	for identification	reage in grain size and exheractial all intermediate particle sixes.	reductionally one size or a range of al with some intermediate sizes aftering.	se or fines with	for identified	fatification Fr	Minteney (Praction to chating)	Outer to alon	Home to very alow	ac ta	Blow to nom	800	Hone to wary alow	milly identified by color, odor, and frequently by fibross texture	itims of group	the minus Po.	Dry Strength (crushing characteristics)	ing particles by or graticles by or graticles by or graticles, the first constitution of the fir
	Tall I		Vide range is a menunis of al	Predominantly of an arternal	Homplantic flam (for identify	Plantic fines (for identification procedure ase CL below).	Vide range is o	Predominantly o	Dominaric fire (for identifi	Plantic from (for see CL below).	on Fraction	Dry Bremgth (Crushing characteristics)	None to elight	Medius to high	Allght to	Slight to	धिक्र क भार संक्र	medium to high	Perdily identify and frequently			Dry Strength (Critical Constitution of the Critical Constitution of the Critical
	tated her		bell-graded gravels, gravel-east airture, little or no fime.	Porly graded grow is or grown-mad mixtures, little or no fines.	filty gravels, gravel-eard-eilt aixture.	Chyey grewle, grewl-sand-clay statures.	well-graded emode, gravelly sends, little or so fisse.	Porty graded made or gravelly made, little or so flass	filty sands, sand-silt sixtures.	Clayey sands, send-clay statures.			Inorganic milts and very fine sands, rock flows, milty or clayer fine sands or clayer milts with milder planticity.	incommute clays of low to medium planticity, gravelly-clays, smally clays, silty clays, less clays.	Organic silts and organic silty clays of low planticity.	Coormanic silts, microsous or distonaceous fine sandy or silty soils, elastic silts.	increase clays of high plasticity, fat clays.	Ormanic clays of medius to high planticity, organic silts.	Past and other highly organic solls.	possessing characteristics of two groups are design	These procedures are to acrest		Little resorting particles larger than Bo, No sieve size, propare a put of moint to still with a volume of about one half citied. And enough moint if moneanty to make he soil out by our not reload. Then the put is the open pain of one hand and the back partition founds of the types of the system to the other hand enough to a larger of the system to the other hand with the other hand has the other hand and the other hand the other hand has the other hand has the other hand the other hand has the other hand to be other hand the other hand to be other hand to be other hand the other hand hand the other hand the other hand hand the other hand had been a with the other hand had been a maken with quits restlict the hand had been a maken with the other hand had been a maken with the hand had been had bee
	3 6	-	в	8	8	R		t _s	ä	8			4	t t	g	•	U	8	ĸ	Bottle year			out one ball to out one ball to out one ball to out one ball to our ball to out out out out out out out out out
	Imper Divisions	2 7	seite in seine sei	nata Prota f Pe sa be	tendo To Mad To	action but and regretarion before the second	AL to la. 1. moles 1. mole 1. mole 1. mole 1. mole 1. mole 2. mole 4. coll mole 5. coll moles 9. coll moles 1. col	of materi	Particle v Band Dark The State State Tresale it attr	and evol sales el	- T V		94 14 P	and tal.	ion ect	eg.	to that a	أساوت	Highly Organic Botla	(1) Boundary classifications:		Dilatancy (reaction to shallag)	After removing particles larger than No. No sizes told that a value of about consult child to the same of about consult child to make the soil north but not without and the large paid of the appearance of where our best mount of the large and best and point the man of the same of the sam

PHOTOGRAPHS OF FROZEN SOIL TYPES



Frozen, well-graded silty SAND. Well-bonded. Classification: SM, Nbn Photograph 3

Frozen lean CLAY. ice lenses in top portion formed from moisture drawn from below. Classification: CL, Vs.r Bottom portion medium bonded and somewhat friable. Classification: CL, Mf

Frozen fine SLLT. Top portion well-bonded, saturated. Classification: ML, Nbn

Photograph 1

Bottom portion friable. Classification: ML, Nf



Photograph 2




Frozen fine SAND. Jell-bonded, high degree of saturation. Classification: SM, Nbe Photograph #


Frozen, clayey sandy GRAVEL with ice coatings on numerous stones. Classification: GW-GC, Vc

Photograph 5



SCALE IN INCHES





SOIL DRY UNIT WEIGHT, ICE VOLUME, AND WATER CONTENT RELATIONSHIPS

ILLUSTRATIVE EXAMPLE OF THE USE OF THE FROZEN SOIL CLASSIFICATION SYSTEM IN TYPICAL EXPLORATION LOG

	0.0		Surface Elevation 963.2 ft
	0.5	OL	Organic SANDY SILT, not frozen.
	0.0	GW	Brown, well-graded SANDY GRAVEL medium compact, moist, not frozen.
	1.8	GW, Nf	Brown, well-graded SANDY GRAVEL, frozen, no visible segregation, negligible thin ice film on gravel sizes and within larger voids, poorly bonded.
	3.7 5.4	GW, Nbn	Brown, well-graded SANDY GRAVEL, frozen, no vicible segregation, well bonded.
	J. 4	ML, Vs	Black, micaceous SANDY SILT, frozen, stratified horizontal ice lenses averaging 4 inches in horizontal extent, hairline to 1/4 inch in thickness, 1/2 to 3/4 inch spacing. Visible ice = 20 ± % of total volume. Ice lenses hard, clear, colorless.
IN FEET	7.7 9.1	ICE	ICE, hard, slightly cloudy, colorless, few scattered inclusions of silty sand.
DEPTH		PT, Vr	Dark brown PEAT, frozen, well bonded, high degree of saturation. 5% visible ice.
	10.5	MH, Vr	Light brown SILT, frozen, irregularly oriented ice lenses and layers $1/4$ to $3/4$ inch thick on random pattern grid approx. 3 to 4 inch spacing. Visible ice = $10\pm\%$ of total volume. Ice moderately soft, porous, gray-white.
	14.3		Bedrock. Laminated shale. Top few feet weathered. I/16" thick ice lenses in fissures to 16.0'. None below.
;	20.6		Bottom of exploration—